

(2) *Sterility*. Proceed as directed in § 436.20 of this chapter, using the method described in paragraph (e)(3) of that section.

(3) *Moisture*. Proceed as directed in § 436.201 of this chapter.

(4) *Metal particles*. Proceed as directed in § 436.206 of this chapter.

[43 FR 11170, Mar. 17, 1978, as amended at 50 FR 19920, May 13, 1985]

**§ 446.381b Tetracycline hydrochloride ophthalmic suspension.**

(a) *Requirements for certification—(1) Standards of identity, strength, quality, and purity*. Tetracycline hydrochloride ophthalmic suspension contains tetracycline hydrochloride in a suitable and harmless oily base. Each milliliter contains 10 milligrams of tetracycline hydrochloride. Its potency is satisfactory if it contains not less than 90 percent and not more than 125 percent of the number of milligrams of tetracycline hydrochloride that it is represented to contain. It is sterile. Its moisture content is not more than 0.5 percent. The tetracycline hydrochloride used conforms to the standards prescribed by § 446.81a(a)(1).

(2) *Labeling*. It shall be labeled in accordance with the requirements of § 432.5 of this chapter.

(3) *Requests for certification; samples*. In addition to complying with the requirements of § 431.1 of this chapter, each such request shall contain:

(i) Results of tests and assays on:

(a) The tetracycline hydrochloride used in making the batch for potency, loss on drying, pH, absorptivity, crystallinity, and identity.

(b) The batch for potency, sterility, and moisture.

(ii) Samples required:

(a) The tetracycline hydrochloride used in making the batch: 10 packages, each containing approximately 300 milligrams.

(b) The batch:

(1) For all tests except sterility: A minimum of five immediate containers.

(2) For sterility testing: 20 immediate containers, collected at regular intervals throughout each filling operation.

(b) *Tests and methods of assay—(1) Potency*. Proceed as directed in § 436.106 of this chapter, preparing the sample for

assay as follows: Place an accurately measured representative portion of the sample into a high-speed glass blender jar with 1 milliliter polysorbate 80 and sufficient 0.1N hydrochloric acid to give a stock solution of convenient concentration containing not less than 150 micrograms of tetracycline hydrochloride per milliliter (estimated). Blend for 3 to 5 minutes. Remove an aliquot and further dilute with sterile distilled water to the reference concentration of 0.24 microgram of tetracycline hydrochloride per milliliter (estimated).

(2) *Sterility*. Proceed as directed in § 436.20 of this chapter, using the method described in paragraph (e)(3) of that section.

(3) *Moisture*. Proceed as directed in § 436.201 of this chapter.

[43 FR 11171, Mar. 17, 1978, as amended at 46 FR 46313, Sept. 18, 1981; 50 FR 19920, May 13, 1985]

**Subpart E—Otic Dosage Forms**

**§ 446.467 Oxytetracycline hydrochloride-polymyxin B sulfate otic ointment.**

(a) *Requirements for certification—(1) Standards of identity, strength, quality, and purity*. Oxytetracycline hydrochloride-polymyxin B sulfate otic ointment is oxytetracycline hydrochloride and polymyxin B sulfate in a suitable and harmless ointment base. Each gram of ointment contains oxytetracycline hydrochloride equivalent to 5 milligrams of oxytetracycline and polymyxin B sulfate equivalent to 10,000 units of polymyxin B. Its oxytetracycline hydrochloride content is satisfactory if it contains not less than 90 percent and not more than 120 percent of the number of milligrams of oxytetracycline that it is represented to contain. Its polymyxin B sulfate content is satisfactory if it contains not less than 90 percent and not more than 125 percent of the number of units of polymyxin B that it is represented to contain. Its moisture content is not more than 1 percent. The oxytetracycline hydrochloride used conforms to the standards prescribed by § 446.67(a)(1). The polymyxin B sulfate used conforms to the standards prescribed by § 448.30(a)(1) of this chapter.